

Total No. of Printed Pages—16

1 SEM TDC GEMT (CBCS) GE 1 (A/B/C)

2 0 2 2

(Nov/Dec)

MATHEMATICS

(Generic Elective)

Paper : GE-1

*The figures in the margin indicate full marks
for the questions*

Paper : GE-1 (A)

(Differential Calculus)

Full Marks : 80

Pass Marks : 32

Time : 3 hours

1. (a) কেতিয়া এটা ফলন f বন্ধ অন্তৰ $[a, b]$ ত অনবচ্ছিন্ন হোৱা বুলি কোৱা হয়? 1

When is a function f said to be continuous in a closed interval $[a, b]$?

- (b) তলৰ যি কোনো এটাৰ মান নিৰ্ণয় কৰা : 3

Evaluate any one of the following :

(i) $\lim_{x \rightarrow 0} \frac{e^x - e^{\sin x}}{x - \sin x}$

(ii) $\lim_{x \rightarrow 0} \frac{\tan x - x}{x - \sin x}$

(2)

(c) f ফলনৰ সংজ্ঞা এনেদৰে দিয়া আছে

$$f(x) = (1+3x)^{1/x}, \quad x \neq 0$$

$$= e^3, \quad x = 0$$

দেখুওৱা যে $x=0$ বিন্দুত ফলন অনবচ্ছিন্ন।

3

Show that the function f defined by

$$f(x) = (1+3x)^{1/x}, \quad x \neq 0$$

$$= e^3, \quad x = 0$$

is continuous at $x=0$.(d) $y = (ax+b)^m$ ৰ n -তম অৱকলজ নিৰ্ণয় কৰা য'ত $n \leq m$ আৰু $m, n \in N$.

1

Find the n -th derivative of $y = (ax+b)^m$, where $n \leq m$ and $m, n \in N$.

(e) যদি (If)

$$y = \frac{\sin^{-1} x}{\sqrt{1-x^2}}$$

দেখুওৱা যে (show that)

$$(1-x^2)y_{n+2} - (2n+3)xy_{n+1} - (n+1)^2 y_n = 0$$

4

2. লিবনিটজৰ উপপাদ্যটো উল্লেখ কৰা আৰু প্ৰমাণ কৰা।

5

State and prove Leibnitz's theorem.

P23/239

(Continued)

(3)

অথবা / Or

যদি (If)

$$u = \tan^{-1} \frac{x^3 + y^3}{x - y}$$

তেজ্জে প্ৰমাণ কৰা যে (then prove that)

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = \sin 2u$$

3. (a) যদি $u = f(xyz)$ হয়, তেজ্জে $\frac{\partial f}{\partial y}$ নিৰ্ণয় কৰা।

1

If $u = f(xyz)$, then find $\frac{\partial f}{\partial y}$.

(b) যদি (If)

$$u = \sin^{-1} \left\{ \frac{\sqrt{x} - \sqrt{y}}{\sqrt{x} + \sqrt{y}} \right\}$$

তেজ্জে প্ৰমাণ কৰা যে (then prove that)

$$\frac{\partial u}{\partial x} = -\frac{y}{x} \frac{\partial u}{\partial y}$$

4

(c) যদি $y = \sin^2 x$, তেজ্জে y_n নিৰ্ণয় কৰা।

1

If $y = \sin^2 x$, then find y_n .

P23/239

(Turn Over)

4. (a) যদি $f = \tan^{-1} \frac{y}{x}$ হয়, তেলে $\frac{\partial f}{\partial x}$ নির্ণয় কৰা।

1

If $f = \tan^{-1} \frac{y}{x}$, then find $\frac{\partial f}{\partial x}$.

(b) দেখুওৱা যে এটা ফলন $f(x) = |x| + |x-1|$, এটা বিন্দু $x=1$ ত অনবচ্ছিন্ন কিন্তু অৱকলনীয় নহয়।

3

Show that the function f defined as follows, is continuous but not derivable at $x=1$, $f(x) = |x| + |x-1|$.

(c) যদি (If)

$$u = \frac{1}{\sqrt{x^2 + y^2 + z^2}}$$

তেলে দেখুওৱা যে (then show that)

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = 0$$

3

5. (a) $y = x^2(a-x)$ বক্রৰ উপস্পর্শকৰ দৈৰ্ঘ্য নির্ণয় কৰা।

1

Find the length of the subtangent to the curve $y = x^2(a-x)$.

(b) দেখুওৱা যে, যি কোনো বক্রৰ ক্ষেত্রত

$$\frac{\text{উপ-অভিলম্ব}}{\text{উপ-স্পর্শক}} = \left(\frac{\text{অভিলম্বৰ দীঘ}}{\text{স্পর্শকৰ দীঘ}} \right)^2$$

2

Show that in any curve

$$\frac{\text{subnormal}}{\text{subtangent}} = \left(\frac{\text{length of normal}}{\text{length of tangent}} \right)^2$$

6. (a) যি কোনো বক্রৰ ক্ষেত্রত উপস্পর্শকৰ সংজ্ঞা লিখা।

1

Define subtangent to any curve.

(b) $x = a(\theta + \sin \theta)$ আৰু $y = a(1 - \cos \theta)$ বক্রৰ θ ত উপস্পর্শকৰ দৈৰ্ঘ্য নির্ণয় কৰা।

3

Find the lengths of subtangent to $x = a(\theta + \sin \theta)$ and $y = a(1 - \cos \theta)$ at θ .

7. তলত দিয়া বক্রৰ অনন্তস্পর্শী নির্ণয় কৰা :

4

Find the asymptotes of the following curve :

$$x^3 - 2x^2y + xy^2 + x^2 - xy + 2 = 0$$

অথবা / Or

$a^4y^2 = x^4(2x^2 - 3a^2)$ বক্রৰ অৱস্থান আৰু দ্বি-বিন্দুৰ প্রকৃতি নির্ণয় কৰা।

Find the position and nature of the double points of the curve $a^4y^2 = x^4(2x^2 - 3a^2)$.

8. তলৰ যি কোনো এটাৰ মান নির্ণয় কৰা :

4

Evaluate any one of the following :

(a) $y = x(x^2 - 1)$ বক্রৰ অনুবেখন নির্ণয় কৰা।

Trace the curve $y = x(x^2 - 1)$.

(b) দেখুওৱা যে $r = a(1 - \cos \theta)$ কাৰ্ডিইডৰ যি কোনো বিন্দু (r, θ) ত বক্রতা ব্যাসার্ধ $\frac{2}{3}\sqrt{2ar}$.

Show that the radius of curvature at any point (r, θ) of the cardioid $r = a(1 - \cos \theta)$ is given by $\frac{2}{3}\sqrt{2ar}$.

9. $f(x, y) = 0$ বক্রৰ যি কোনো বিন্দু (x, y) ত বহু বিন্দু হোৱাৰ প্ৰয়োজনীয় আৰু পৰ্যাপ্ত চৰ্ত উল্লেখ কৰি প্ৰমাণ কৰা। 5

State and prove the necessary and sufficient condition for any point (x, y) on the curve $f(x, y) = 0$ to be a multiple point.

অথবা / Or

এটা বক্রৰ কাৰ্টেচিয়ান সমীকৰণ $y = f(x)$ হ'লে বক্রৰ এটা বিন্দুত বক্রতা ব্যাসাৰ্থ নিৰ্ণয় কৰা।

Find the radius of curvature at a point of the Cartesian equation of the curve $y = f(x)$.

10. (a) ৰোলৰ উপপাদ্যটো লিখা। 1

State the Rolle's theorem.

- (b) $[-1, 1]$ অন্তৰালত $f(x) = \frac{1}{2-x^2}$ ফলনৰ বাবে ৰোলৰ

উপপাদ্য প্ৰতিপন্ন কৰা। 2

Verify Rolle's theorem for the function

$$f(x) = \frac{1}{2-x^2}$$

in the interval $[-1, 1]$.

- (c) মধ্যমান উপপাদ্য $f(b) - f(a) = (b-a)f'(\xi)$ প্ৰতিপন্ন কৰা য'ত $f(x) = x(x-1)(x-3)$, $a=0$, $b=\frac{1}{2}$

আৰু ξ ব মান নিৰ্ণয় কৰা। 4

Verify the applicability of the mean value theorem $f(b) - f(a) = (b-a)f'(\xi)$, $a < \xi < b$ if $f(x) = x(x-1)(x-3)$, where $a=0$, $b=\frac{1}{2}$. Also find the value of ξ .

11. লাগ্ৰাঞ্জৰ মধ্যমান উপপাদ্য উল্লেখ কৰি প্ৰমাণ কৰা। $1+4=5$

State and prove Lagrange's mean value theorem.

অথবা / Or

মেক্লেৰিনৰ উপপাদ্য ব্যৱহাৰ কৰি $\sin x$ ক x -ৰ সূচকত অসীম শ্ৰেণীত বিস্তৃতি কৰা। 5

Using Maclaurin's theorem, expand $\sin x$ in an infinite series in powers of x .

12. (a) যদি (If)

$$f(x) = f(0) + xf'(0) + \frac{x^2}{2} f''(\theta x)$$

তেতিয়া θ ব মান উলিওৱা যেতিয়া $x \rightarrow 1$ আৰু য'ত $f(x) = (1-x)^{5/2}$.

then find θ when $x \rightarrow 1$ and where $f(x) = (1-x)^{5/2}$. 3

- (b) $f(x, y) = x^3 + y^3 - 3x - 12x + 20$ ফলনৰ সৰ্বোচ্চ আৰু সৰ্বনিম্ন মান নিৰ্ণয় কৰা। 4

Find the maximum and minimum values of the function

$$f(x, y) = x^3 + y^3 - 3x - 12x + 20$$

13. (a) $\log x$ ক $x-1$ ব সূচকত বিস্তৃতি কৰা য'ত
 $0 < x \leq 2$. 1
 Expand $\log x$ in powers of $x-1$ where
 $0 < x \leq 2$.

- (b) তলৰ যি কোনো এটাৰ মান নিৰ্ণয় কৰা : 4
 Evaluate any one of the following :

$$(i) \lim_{x \rightarrow 1} \left\{ \frac{x}{x-1} - \frac{1}{\log x} \right\}$$

$$(ii) \lim_{x \rightarrow 0} (\cos x)^{\cot^2 x}$$

14. (a) লাগ্ৰাঞ্জৰ ৰূপৰ অৱশেষ থকা মেক্লেৰিনৰ উপপাদ্য লিখা। 1
 Write the Maclaurin's theorem with
 Lagrange's form of remainder.

- (b) মেক্লেৰিনৰ অসীম শ্ৰেণী ব্যৱহাৰ কৰি $\log(1+x)$ ৰ
 বিস্তৃতি কৰা য'ত $-1 < x < 1$. 5
 Expand $\log(1+x)$ using Maclaurin's
 infinite series where $-1 < x < 1$.

অথবা / Or

লাগ্ৰাঞ্জৰ ৰূপৰ অৱশেষ থকা টেইলৰ উপপাদ্য লিখি
 প্ৰমাণ কৰা।

State and prove Taylor's theorem with
 Lagrange's form of remainder.

Paper : GE-1 (B)

(Object-Oriented Programming in C++)

Full Marks : 60

Pass Marks : 24

Time : 3 hours

1. Answer the following questions : 1×5=5

- (a) Define abstraction.
 (b) State one difference between C and C++.
 (c) Write one characteristic of object-oriented programming language.
 (d) What is the use of <iostream.h>?
 (e) How are objects created from a class?

2. Answer any five of the following questions :

2×5=10

- (a) When do you declare a method or class abstract?
 (b) Briefly explain the structure of C++ program.
 (c) How does inheritance help us to create new classes?
 (d) Why can we not override static method?

- (e) State the difference between while loop and do while loop.
- (f) Define default constructor and copy constructor.

3. Answer any *five* of the following questions :

3×5=15

- (a) Explain the following operators and their uses :
cin, cout and delete.
- (b) Explain the three access modifiers.
- (c) What is dynamic binding? Define message passing.
- (d) State the difference between break and continue with example.
- (e) Define file pointer. What is function prototyping? Explain with example.
- (f) Explain the increment and decrement operators in brief.

4. Answer any *four* of the following questions :

5×4=20

- (a) Write a C++ program to store information of a book in a structure.
- (b) Write a C++ program to overload a unary operator.

- (c) Write a C++ program to display Fibonacci series up to 50.
- (d) Write a C++ program to implement friend function.
- (e) Write a C++ program to count the number of objects created.

5. (a) Explain the different types of inheritance with examples and diagrams.

10

Or

- (b) Explain inline and virtual functions with suitable example.

Paper : GE-1 (C)

(Finite Element Methods)

Full Marks : 80

Pass Marks : 32

Time : 3 hours

1. (a) Write True or False : 1
The finite-element method is a piecewise application of a variational method.
- (b) Write down the differences between finite difference methods and finite element methods. 3
- (c) Consider the boundary value problem

$$u'' + (1+x^2)u + 1 = 0$$
Determine the coefficients of the approximate solution

$$W(x) = a_1(1-x^2) + a_2x^2(1-x^2)$$
by using the least square method. 5
- Or
- Using Galerkin's method, solve the boundary value problem

$$\nabla^2 u = -1, \quad |x| \leq 1, \quad |y| \leq 1$$

$$u = 0, \quad |x| = 1, \quad |y| = 1$$
with $h = \frac{1}{2}$.

- (d) Find the variational functional for the boundary value problem

$$u'' = u - 4xe^x$$

$$u'(0) - u(0) = 1, \quad u'(1) + u(1) = -e$$
 5
- (e) State and prove the Lax-Milgram theorem. 6
2. (a) The application of the finite element method to the boundary value problem

$$-u'' = x$$

$$u(0) = u(1) = 0$$
leads to the system of equations $Au = b$. Determine the matrix A and the column vector b for four elements of equal lengths. 6
- (b) Apply Galerkin method to the boundary value problem

$$\nabla^2 u + \lambda u = 0, \quad |x| \leq 1, \quad |y| \leq 1$$

$$u = 0, \quad |x| = 1, \quad |y| = 1$$
to get the characteristic equation in the form $|A - \lambda B| = 0$. 6
3. (a) Define assembly of the element equations. 1
- (b) Define two principles that were used in one-dimensional problem to assembly of finite element equations. 2

- (c) Discuss briefly with an example about the element assemblage in finite element method. 3
- (d) Write down the importance of sparse matrix in the process of element assemblage with an example. 4
- (e) If the finite solutions at any point (x, y) in an element Ω^e is given by


$$U(x, y) = \sum_{J=1}^n U_J^e \psi_J^e(x, y)$$

Find its derivatives. 2

- 4. (a) State the properties for a quadratic triangular element. 3
- (b) Give an example of triangular element with a common node. 1
- (c) Illustrate the process of discretization in two-dimensional domain with a suitable example. 5
- (d) Write the importance of isoperimetric element in the process of element assemblage with an example. 3

- 5. (a) Write True or False : 1
 Finite element modelling involves assumptions concerning the representation of the system and its behaviour.

- (b) Write about interpolating function in finite element method. Find an expression for interpolating function in one-dimensional domain. 3
- (c) Calculate the interpolation function for the quadratic triangular element shown in the figure : 4




- (d) Evaluate the integral of the form

$$I = \int_{(e)} F(x) dx$$

for the triangular element where $F(x)$ is given function, (e) is the element and x represents multidimensional coordinates. 4

Or

Consider the quadratic triangular element shown in the figure :



Evaluate the integral of the product

$$\left(\frac{\partial \psi_1}{\partial x} \right) \left(\frac{\partial \psi_4}{\partial x} \right)$$

at the point $(x, y) = (2, 4)$.

6. (a) What are the different types of partial differential equations? Write their field in applications. 4

- (b) Find the solution of the boundary value problem

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{1+e^u}{2} = 0, \quad |x| \leq 1, \quad |y| \leq 1$$
$$u = 0, \quad |x| = 1, \quad |y| = 1$$

by finite element method (use the three node triangular element). 4

- (c) Use finite element method to solve the boundary value problem

$$\nabla^2 u = -1, \quad |x| \leq 1, \quad |y| \leq 1$$
$$\frac{\partial u}{\partial x} + u = 0, \quad |x| = 1, \quad |y| = 1$$

with $h = \frac{1}{2}$. 4
