

Total No. of Printed Pages—6

6 SEM TDC DSE PHY (CBCS) 1 (H)

2 0 2 3

(May/June)

PHYSICS

(Discipline Specific Elective)

(For Honours)

Paper : DSE-1

(Nuclear and Particle Physics)

Full Marks : 80

Pass Marks : 32

Time : 3 hours

*The figures in the margin indicate full marks
for the questions*

1. Choose the correct option from the following :

1×5=5

(a) With increase in mass number, the
neutron to proton ratio

(i) increases

(ii) decreases

(iii) increases first and then decreases

(iv) None of the above

(2)

- (b) Neutrons have a _____ value of dipole magnetic moment.
- (i) positive
 - (ii) negative
 - (iii) zero
 - (iv) None of the above
- (c) In alpha decay
- (i) mass number A decreases by 4 and atomic number Z increases by 2
 - (ii) mass number A decreases by 4 and atomic number Z decreases by 2
 - (iii) mass number A increases by 4 and atomic number Z decreases by 2
 - (iv) mass number A increases by 4 and atomic number Z increases by 2
- (d) Electron is a _____ generation particle.
- (i) first
 - (ii) second
 - (iii) third
 - (iv) None of the above
- (e) Isospin is to be conserved in
- (i) all elementary interactions
 - (ii) strong interactions only
 - (iii) weak interactions only
 - (iv) None of the above

P23/824

(Continued)

(3)

2. (a) What is nuclear quadrupole moment? 1
- (b) Write down the relation between mass number and radius of a nucleus. Describe a method for determining nuclear radius. 1+3=4
- Or
- Explain the terms 'nuclear angular momentum' and 'nuclear quadrupole moment'. 2+2=4
3. (a) What are magic numbers? What is their significance in the shell model of the nucleus? 1+2=3
- (b) What are the applications of the semi-empirical mass formula? Draw a graph indicating the contribution of the various terms of the semi-empirical mass formula to the total binding energy. 2+2=4
- (c) Describe the liquid-drop model of the nucleus describing the similarities of the nucleus with a drop of liquid. How can nuclear fission be explained on the basis of this model? 4+2=6

P23/824

(Turn Over)

(4)

4. (a) Describe how the range of alpha particles can be determined. What is straggling? Write down the relation connecting range and disintegration constant. $3+1+1=5$
- (b) Write down the equation showing the three modes of beta radioactivity. Describe the role of neutrino in explaining continuous energy spectrum of beta particles. $1+3=4$
5. (a) Discuss Rutherford scattering in a nucleus. 3
- (b) Derive an expression for Q-value of a nuclear reaction. 3
- (c) A 7.7 MeV alpha particle interacts with a target nucleus $^{14}_7\text{N}$ to produce a residual nucleus $^{17}_8\text{N}$ and a product particle ^1_1H . The protons emitted at 90° to the incident beam direction are found to have kinetic energy of 4.44 MeV. Calculate the Q-value of the reaction. 3

Or

Differentiate between direct reaction and compound nucleus reaction.

P23/824

(Continued)

(5)

6. Write short notes on any two of the following : $4 \times 2 = 8$
- (a) Cerenkov radiation
- (b) Photoelectric effect
- (c) Interaction of neutron with matter
7. What are the gas filled detectors? Describe briefly how gas filled detectors work in the following different regions on varying the plate voltage : $1+2+3+3=9$
- (a) Ionization chamber region
- (b) Proportional region
- (c) Geiger region

Or

Describe the principle and working of a scintillation detector. Name any two scintillators. Describe the working of a photomultiplier tube. $4+2+3=9$

8. Describe the working of a cyclotron. How are the difficulties faced in a cyclotron removed in a synchrotron? $3+1+1=5$

Or

What are tandem accelerators? Describe the construction and working of a van de Graaff generator. $2+3=5$

P23/824

(Turn Over)

9. (a) What is strong interaction? What are the conservation laws to be satisfied in strong interaction? What is the associated exchange particle? $1+2+1=4$
- (b) What does generation mean in particle physics? Which particles are the first-, second- and third-generation leptons? $1+3=4$
- (c) What are hadrons? Which fundamental interaction is specific to them? $2+1=3$
- (d) Check whether isospin and strangeness are conserved in the following reactions : $2 \times 3 = 6$
- (i) $\pi^+ + n \rightarrow \pi^- + p$
- (ii) $\pi^- + p \rightarrow \Lambda^0 + K^0$
- (iii) $\pi^+ + \Lambda^0 \rightarrow \Sigma^+ + K^0$

Or

What are quarks? Give the quark structure of pions. $3+3=6$

★ ★ ★