4 SEM TDC PHYH (CBCS) C 8

2023

(May/June)

PHYSICS

(Core)

Paper: C-8

(Mathematical Physics—III)

Full Marks: 53
Pass Marks: 21

Time: 3 hours

The figures in the margin indicate full marks for the questions

1. Choose the correct answer:

 $1 \times 4 = 4$

- (a) The value of $|3z_1 4z_2|$, if $z_1 = 2 + i$ and $z_2 = 3 2i$, is
 - (i) 157
 - (ii) 167
 - (iii) √157
 - (iv) √167

- (b) The value of the integral $\oint_C \frac{1}{z} dz$, if the point z = 0 lies within the curve C, is
 - (i) $\frac{1}{2\pi i}$
 - (ii) O
 - (iii) 2πi
 - (iv) None of the above
- (c) The Laplace transform f(s) of F(t) = t is
 - (i) 1
 - (ii) s
 - (iii) s2
 - (iv) $1/s^2$
- (d) If $g(\omega)$ is the Fourier transform of f(t), the Fourier transform of f(at) is
 - (i) $\frac{1}{a}g\left(\frac{\omega}{a}\right)$
 - (ii) $\frac{1}{\omega}g\left(\frac{\omega}{a}\right)$
 - (iii) $\frac{1}{\omega}g\left(\frac{a}{\omega}\right)$
 - (iv) None of the above

2. Answer the following:

2×5=10

- (a) Express the equation $x^2 + y^2 = 36$ in terms of the conjugate co-ordinates -5 + 5i.
- (b) Express the complex number $-1 + \sqrt{3}i$ in polar form.
- (c) Differentiate between the terms 'essential singularity' and 'removable singularity' with examples.
- (d) Using the Fourier integral formula, derive the Fourier cosine transformation.
- (e) Prove that if f(s) is the Laplace transform of F(t), the Laplace transform of $e^{at}F(t)$ is f(s-a).
- 3. Answer the following:
 - (a) State the Cauchy's residue theorem. Prove that $\int_0^\infty \frac{dx}{x^6 + 1} = \frac{\pi}{3}$. 1+4=5
 - (b) If f(z) = u + iv is analytic and $u = e^{-x}(x \sin y y \cos y)$, prove that u is harmonic. Also, find the value of v. 2+3=5
 - (c) State the Cauchy's integral formula for derivative of a function. Evaluate the integral $\oint_C \frac{e^{2z}}{(z+1)^4} dz$, where C is the circle |z| = 3.

(d) Find the residues of vollal and toward .

$$f(z) = \frac{z^2 - 2z}{(z+1)^2(z^2+4)}$$

3

- (e) Find the Laurent series of $\frac{e^{2z}}{(z-1)^3}$ about z=1.
- **4.** Find the Fourier transform of the following functions (any *two*): $3\times2=6$

(i)
$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right)$$

(ii) $f(x) = Ne^{-\alpha x^2}$ (N and α are constants)

(iii)
$$f(x) = \frac{1}{2\pi} \left(\frac{a^2}{x^2 + a^2} \right)$$

- **5.** Find the Laplace transform of the following functions (any two): $3\times2=6$
 - (i) $f(t) = \frac{e^{at} 1}{a}$
 - (ii) $f(t) = \cos^2 t$
 - (iii) f(t) = Kt, where K is a constant
- **6.** Write short notes on the following (any *two*): $3\times 2=6$
 - (a) Singularities of a complex function
 - (b) Fourier transforms and its applications
 - (c) Convolution theorem

* * *